Two-parameter Quantum Affine Superalgebra $U_{p,q}(gl\widehat{(1|1)})$ and Its Drinfel′d's Realization

^{a}Jin-fang Cai[1] , ^{bc}Shi-kun Wang, ^{a}Ke Wu

^{a} Institute of Theoretical Physics, Academia Sinica,
Beijing, 100080, P. R. China

^{b} CCAST (World Laboratory), P.O. Box 3730, Beijing, 100080, P. R. China

^{c} Institute of Applied Mathematics, Academia Sinica,
Beijing, 100080, P. R. China

arXiv:q-alg/9703030v1 17 Mar 1997

Abstract

Using super RS construction method and Gauss decomposition, we obtain Drinfel′d's currents realization of two-parameter quantum affine superalgebra $U_{p,q}(gl\widehat{(1|1)})$ and get co-product structure for these currents

For quantum affine algebras, there are three realizations or constructions: Drinfel′d-Jimbo realization [1, 2], Drinfel′d [3] and RS [4] (or affine version of FRT [5]) realization. The explicit isomorphism between Drinfel′d and RS realization was established by Ding and Frenkel [6] using Gauss decomposition. Recently, some attensions have been paid to the construction or definition of quantum affine superalgebras [7, 8, 9]. In this paper, we will use the super version of RS construction method and Ding-Frenkel isomorphic map to define a two-parameter quantum affine superalgebra $U_{p,q}(gl\widehat{(1|1)})$ and get its Drinfel′d's currents realization. We also calculate co-product structure for these currents.

The graded Yang-Baxter equation(YBE) takes this form [10]:

$$\eta_{12}R_{12}(z/w)\eta_{13}R_{13}(z)\eta_{23}R_{23}(w) = \eta_{23}R_{23}(w)\eta_{13}R_{13}(z)\eta_{12}R_{12}(z/w), \tag{1}$$

where R-matrix acts in tensor product of two 2-D graded linear space V: $R(z) \in End(V \otimes V)$, and $\eta_{ik,jl} = (-1)^{(i-1)(k-1)}\delta_{ij}\delta_{lk}$. It can also be writen in components as:

$$\begin{aligned} &R^{ij}_{ab}(z/w)R^{ak}_{pc}(z)R^{bc}_{qr}(w)(-1)^{(P(a)-P(p))P(b)} \\ &\qquad = (-1)^{P(e)(P(f)-P(r))}R^{jk}_{ef}(w)R^{if}_{dr}(z)R^{de}_{pq}(z/w) \end{aligned} \tag{2}$$

It's very obvious that $\eta R(z)$ satisfies the ordinary YBE when $R(z)$ is a solution of graded

---

[1] *Email address: caijf@itp.ac.cn*

YBE. It can be verified that following R-matrix [10] is a solution of graded YBE (1) :

$$R_{12}(z) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \frac{(z-1)qp^{-1}}{zq-p^{-1}} & \frac{z(q-p^{-1})}{zq-p^{-1}} & 0 \\ 0 & \frac{(q-p^{-1})}{zq-p^{-1}} & \frac{z-1}{zq-p^{-1}} & 0 \\ 0 & 0 & 0 & -\frac{q-zp^{-1}}{zq-p^{-1}} \end{pmatrix} \tag{3}$$

This solution satisfy the unitary condition: $R_{12}(z)R_{21}(z^{-1}) = \mathbf{1}$, and it have two deformation parameters: p and q. From the above solution of graded YBE, we can define two-parameter quantum affine superalgebra $U_{p,q}(gl\widehat{(1|1)})$ with a central extension employing the super version RS construction method [4]. $U_{p,q}(gl\widehat{(1|1)})$ is an associative algebra with generators $\{l^k_{ij}|1 \leq i,j \leq 2, k \in \mathbf{Z}\}$ which subject to the following multiplication relations:

$$R_{12}(\tfrac{z}{w})L^\pm_1(z)\eta L^\pm_2(w)\eta = \eta L^\pm_2(w)\eta L^\pm_1(z)R_{12}(\tfrac{z}{w}) \tag{4}$$

$$R_{12}(\tfrac{z_-}{w_+})L^+_1(z)\eta L^-_2(w)\eta = \eta L^-_2(w)\eta L^+_1(z)R_{12}(\tfrac{z_+}{w_-}) \tag{5}$$

here $z_\pm = zq^{\pm\frac{c}{2}}$. We have used standard notation: $L^\pm_1(z) = L^\pm(z)\otimes\mathbf{1}$, $L^\pm_2(z) = \mathbf{1}\otimes L^\pm(z)$ and $L^\pm(z) = \left(l^\pm_{ij}(z)\right)^2_{i,j=1}$, $l^\pm_{ij}(z)$ are generating functions (or currents) of l^k_{ij}: $l^\pm_{ij}(z) = \sum_{k=0}^\infty l^{\pm k}_{ij} z^{\pm k}$.

This algebra admits coalgebra and antipole structure compatible with the associative multiplication defined by eqs.(4)(5):

$$\triangle\left(l^\pm_{ij}(z)\right) = \sum_{k=1}^{2} l^\pm_{kj}(zq^{\pm\frac{c_2}{2}})\otimes l^\pm_{ik}(zq^{\mp\frac{c_1}{2}})(-1)^{(k+i)(k+j)}, \tag{6}$$

$$\epsilon\left(l^\pm_{ij}(z)\right) = \delta_{ij}, \qquad S\left({}^{st}L^\pm(z)\right) = \left[{}^{st}L^\pm(z)\right]^{-1}. \tag{7}$$

where $\left[{}^{st}L^\pm(z)\right]_{ij} = (-1)^{i+j}l^\pm_{ji}(z)$.

$L^\pm(z)$ have the following unique decompositions [6] :

$$L^\pm(z) \;=\; \begin{pmatrix} 1 & 0 \\ f^\pm(z) & 1 \end{pmatrix}\begin{pmatrix} k^\pm_1(z) & 0 \\ 0 & k^\pm_2(z) \end{pmatrix}\begin{pmatrix} 1 & e^\pm(z) \\ 0 & 1 \end{pmatrix} \tag{8}$$

where $e^\pm(z)$, $f^\pm(z)$ and $k^\pm_i(z)$ (i=1,2) is generating functions of $U_{p,q}(gl\widehat{(1|1)})$ and $k^\pm_i(z)$ (i=1,2) are invertible. Let $X^+(z) = e^+(z_-) - e^-(z_+)$ and $X^-(z) = f^+(z_+) - f^-(z_-)$. By the similar calculation made in [6, 8] for quantum affine (super) algebras, we obtain the following (anti-)commutation relations among $X^\pm(z)$ and $k^\pm_i(z)$ (i=1,2):

$$[k^\pm_1(z)\ ,\ k^\pm_1(w)] = [k^+_1(z)\ ,\ k^-_1(w)] = 0 \tag{9}$$

$$[k^\pm_1(z)\ ,\ k^\pm_2(w)] = [k^\pm_2(z)\ ,\ k^\pm_2(w)] = 0 \tag{10}$$

$$\frac{z_\pm - w_\mp}{z_\pm q - w_\mp p^{-1}}k^\pm_1(z)k^\mp_2(w)^{-1} = \frac{z_\mp - w_\pm}{z_\mp q - w_\pm p^{-1}}k^\mp_2(w)^{-1}k^\pm_1(z) \tag{11}$$

$$\frac{w_+q - z_-p^{-1}}{z_-q - w_+p^{-1}}k^+_2(z)^{-1}k^-_2(w)^{-1} = \frac{w_-q - z_+p^{-1}}{z_+q - w_-p^{-1}}k^-_2(w)^{-1}k^+_2(z)^{-1} \tag{12}$$

$$k_i^{\pm}(z)^{-1}X^{+}(w)k_i^{\pm}(z)=\frac{z_{\pm}p-wq^{-1}}{z_{\pm}-w}X^{+}(w)\qquad(i=1,2)\tag{13}$$

$$k_i^{\pm}(z)X^{-}(w)k_i^{\pm}(z)^{-1}=\frac{z_{\mp}p-wq^{-1}}{z_{\mp}-w}X^{-}(w)\qquad(i=1,2)\tag{14}$$

$$\{X^{+}(z)\ ,\ X^{+}(w)\}=\{X^{-}(z)\ ,\ X^{-}(w)\}=0\tag{15}$$

$$\{X^{+}(z)\ ,\ X^{-}(w)\}=(p-q^{-1})\left[\delta(\frac{w_{-}}{z_{+}})k_1^{-}(z_{+})^{-1}k_2^{-}(z_{+})-\delta(\frac{z_{-}}{w_{+}})k_1^{+}(w_{+})^{-1}k_2^{+}(w_{+})\right]\tag{16}$$

here $\delta(z)=\sum_{k\in\mathbf{Z}}z^k$. The above relations are Drinfel′d's currents realization of two-parameter quantum affine superalgebra $U_{p,q}(gl\widehat{(1|1)})$. It's very clear that $X^{\pm}(z)$ (or $e^{\pm}(z)$ and $f^{\pm}(z)$) are Fermionic type for their anti-commutation relations and $k_i^{\pm}(z)$ (i=1,2) are Bosonic type elements in $U_{p,q}(gl\widehat{(1|1)})$ as expected.

Introducing a transformation for currents:

$$E(z)=X^{+}(zq)\qquad F(z)=X^{-}(zq)\tag{17}$$

$$K^{\pm}(z)=k_1^{\pm}(zq)^{-1}k_2^{\pm}(zq)\qquad H^{\pm}(z)=k_2^{\pm}(zq)k_1^{\pm}(zp^{-1})\tag{18}$$

then the (anti-)commutation relations for $E(z),F(z),K^{\pm}(z)$ and $H^{\pm}(z)$ are:

$$[K^{\pm}(z)\ ,\ K^{\pm}(w)]=[H^{\pm}(z)\ ,\ H^{\pm}(w)]=0\tag{19}$$

$$[K^{+}(z)\ ,\ K^{-}(w)]=[K^{\pm}(z)\ ,\ H^{\pm}(w)]=0\tag{20}$$

$$\frac{w_{\pm}q-z_{\mp}p^{-1}}{w_{\pm}q^{-1}-z_{\mp}p}K^{\pm}(z)H^{\mp}(w)=H^{\mp}(w)K^{\pm}(z)\frac{w_{\mp}q-z_{\pm}p^{-1}}{w_{\mp}q^{-1}-z_{\pm}p}\tag{21}$$

$$\left(\frac{z_{+}q-w_{-}p^{-1}}{z_{+}p^{-1}-w_{-}q}\right)^2H^{+}(z)H^{-}(w)=H^{-}(w)H^{+}(z)\left(\frac{z_{-}q-w_{+}p^{-1}}{z_{-}p^{-1}-w_{+}q}\right)^2\tag{22}$$

$$[K^{\pm}(z)\ ,\ E(w)]=[K^{\pm}(z)\ ,\ F(w)]=0\tag{23}$$

$$E(w)H^{\pm}(z)=\frac{z_{\pm}p-wq^{-1}}{z_{\pm}p^{-1}-wq}H^{\pm}(z)E(w)\tag{24}$$

$$H^{\pm}(z)F(w)=\frac{z_{\mp}p-wq^{-1}}{z_{\mp}p^{-1}-wq}F(w)H^{\pm}(z)\tag{25}$$

$$\{E(z)\ ,\ E(w)\}=\{F(z)\ ,\ F(w)\}=0\tag{26}$$

$$\{E(z)\ ,\ F(w)\}=(p-q^{-1})\left[\delta(\frac{w_{-}}{z_{+}})K^{-}(z_{+})-\delta(\frac{z_{-}}{w_{+}})K^{+}(w_{+})\right]\tag{27}$$

The co-product structure for currents can be calculated directly from (6)(7):

$$\triangle\left(K^{\pm}(z)\right)=K^{\pm}(zq^{\pm\frac{c_2}{2}})\otimes K^{\pm}(zq^{\mp\frac{c_1}{2}})\tag{28}$$

$$\begin{aligned}\triangle\left(H^{\pm}(z)\right)=&H^{\pm}(zq^{\pm\frac{c_2}{2}})\otimes H^{\pm}(zq^{\mp\frac{c_1}{2}})-(p+q^{-1})F^{\pm}(zp^{-1}q^{-1\mp\frac{c_1}{2}\pm\frac{c_2}{2}})H^{\pm}(zq^{\pm\frac{c_2}{2}})\otimes\\&H^{\pm}(zq^{\mp\frac{c_1}{2}})E^{\pm}(zp^{-1}q^{-1\mp\frac{c_1}{2}\pm\frac{c_2}{2}})\end{aligned}\tag{29}$$

$$\triangle\left(E^{\pm}(z)\right)=E^{\pm}(z)\otimes\mathbf{1}+K^{\pm}(zq^{\mp\frac{c_1}{2}})\otimes E^{\pm}(zq^{\mp c_1})\tag{30}$$

$$\triangle\left(F^{\pm}(z)\right)=\mathbf{1}\otimes F^{\pm}(z)+F^{\pm}(zq^{\pm c_2})\otimes K^{\pm}(zq^{\pm\frac{c_2}{2}})\tag{31}$$

$$\epsilon\left(K^{\pm}(z)\right)=\epsilon\left(K^{\pm}(z)\right)=1\tag{32}$$

$$\epsilon\left(E^{\pm}(z)\right)=\epsilon\left(F^{\pm}(z)\right)=0\tag{33}$$

where $E^{\pm}(z) = e^{\pm}(z_{\mp}q)$ and $F^{\pm}(z) = f^{\pm}(z_{\pm}q)$, then $E(z) = E^{+}(z) - E^{-}(z)$ and $F(z) = F^{+}(z) - F^{-}(z)$.

References

[1] V.G. Drinfel′d, "Quantum Groups", *Proc.ICM-86(Berkeley)*. Vol.1. NewYork Academic Press 1986, 789

[2] M. Jimbo, *Commun. Math. Phys.* **102** (1986) 537

[3] V.G. Drinfel′d: *Soviet Math. Dokl.* **36** (1988) 212

[4] N.Yu. Reshetikhin and M.A. Semenov-Tian-Shansky, *Lett. Math. Phys.* **19** (1990) 133

[5] L.D. Faddeev, N.Yu. Reshetikhin and L.A. Takhtajan, *Algebraic analysis* **1** (1988) 129

[6] J. Ding and I.B. Frenkel, *Commun. Math. Phys.* **156** (1993) 277

[7] H. Yamane, "On Definding Relations Of The Affine Lie Superalgebras And Their Quantized Universal Enveloping Superalgebra", preprint, q-alg/**9603015**

[8] J.F. Cai, S.K. Wang and K. Wu and W.Z. Zhao, "Drinfel′d Realization of Quantum Affine Superalgebra $U_q(gl\widehat{(1|1)})$", preprint, q-alg/**9703022**

[9] Y.Z. Zhang, "Comments on Drinfeld Realization of Quantum Affine Superalgebra $U_q[gl(m|n)^{(1)}]$ and Its Hopf Algebra Structure", preprint, q-alg/**9703020**

[10] H. Yan, Y. Zhou and T.H. Zhu, *J. Phys.* **A26** (1993) 935